

Cephalopod paralarvae assemblages in Hawaiian Islands waters

John R. Bower^{1,*}, Michael P. Seki², Richard E. Young³, Keith A. Bigelow⁴,
Jed Hirota³, Pierre Flament³

¹Faculty of Fisheries, Hokkaido University, 3-1-1 Minato-cho, Hakodate, Hokkaido, 041-8611, Japan

²Honolulu Laboratory, Southwest Fisheries Science Center, National Marine Fisheries Service, NOAA, 2570 Dole Street, Honolulu, Hawaii 96822-2396, USA

³Department of Oceanography, University of Hawaii at Manoa, 1000 Pope Road, Honolulu, Hawaii 96822, USA

⁴Oceanic Fisheries Programme, Secretariat of the Pacific Community, BP D5, 98848 Noumea Cedex, New Caledonia

ABSTRACT: The distribution and abundance of cephalopod paralarvae near the Hawaiian Islands are described. Paralarvae were collected during 5 plankton surveys in 1991 to 1993. The 404 tows at 59 stations collected 10375 paralarvae from 21 families and 57 species. The most numerous families were the Ommastrephidae (23% of total catch), Pyroteuthidae (17%), Enoploteuthidae (16%), Onychoteuthidae (14%), and Ctenopterygidae (8%). The most numerous species were *Ommastrephes bartramii* (18%), *Pterygioteuthis microlampas* (15%), *Ctenopteryx sicula* (8%), and *Onychoteuthis compacta* (6%). Analysis of paralarval distribution patterns identified 2 paralarval assemblages: 'island associated' and 'oceanic'. The 15 'island-associated' species showed increased paralarval abundance near the islands, suggesting preferential spawning in this area. Epipelagic nearshore spawners included *Onychoteuthis* sp. C, *Sthenoteuthis oualaniensis*, and *Nototodarus hawaiiensis*. Mesopelagic nearshore spawners included 3 reported members of the Hawaiian Mesopelagic Boundary Community (*Abralia trigonura*, *Liocranchia reinhardti*, and *Chiroteuthis picteti*) and 3 probable new members (*Liocranchia valdiviae*, *Histioteuthis hoylei* and *Enoploteuthis jonesi*).

KEY WORDS: Paralarvae · Cephalopod · Hawaiian Islands · Distribution · Assemblage

INTRODUCTION

Cephalopod paralarvae are more numerous and more easily sampled than adults. Knowledge of paralarval distribution and abundance patterns is useful for determining when and where adults spawn, particularly for species whose adults are difficult to catch. Knowledge of the early life stages can also help in understanding cephalopod population dynamics and in developing stock-recruitment models for commercially important species (Vecchione 1987). Yet, despite their importance, the early life stages of most species are seldom studied and poorly understood (Boyle 1990).

A diverse community of cephalopods occurs near the Hawaiian Islands (Berry 1914, Young 1978). It includes

inhabitants of the mesopelagic-boundary region (Reid et al. 1991, Young 1995), one of the most poorly investigated habitats in the ocean. Benthic adults are particularly difficult to sample because the steep and jagged seafloor near the islands restricts benthic trawling. As a result, the species composition and distribution of cephalopods around the islands remain uncertain.

To date, the paralarval distribution patterns of only 1 cephalopod species (*Ommastrephes bartramii*) near the Hawaiian Islands have been described (Young & Hirota 1990, Bower 1994, 1996). In this paper, we present the paralarval distribution patterns of 58 cephalopod species collected near the islands during 5 surveys in 1991 to 1993. We then identify species-assemblage groups based on capture distance from the islands and compare onshore-offshore abundance patterns to infer which species may spawn preferentially near the islands.

*E-mail: akaika@pop.fish.hokudai.ac.jp

MATERIALS AND METHODS

Sampling. Five plankton surveys were conducted near the Hawaiian Islands to collect cephalopod paralarvae. Stations were located 9 to 1169 km windward (northeast) and 11 to 256 km leeward (southwest) of the islands. Samples were collected aboard the Hokkaido University ship 'Hokusei Maru' (HM) during 6 to 15 February 1991 (HM-91 survey; 7 stations, 3 to 5 tows station⁻¹), 5 to 26 February 1992 (HM-92 survey; 16 stations, 7 to 10 tows station⁻¹), and 4 to 19 February 1993 (HM-93 survey; 12 stations, 7 to 10 tows station⁻¹), and aboard the NOAA ship 'Townsend Cromwell' (TC) during 5 to 20 February 1991 (TC-91 survey; 10 stations, 1 to 7 tows station⁻¹) and 22 March to 7 April 1992 (TC-92 survey; 14 stations, 4 to 8 tows station⁻¹). The standard sampling procedure in all surveys was to conduct 30 min open oblique tows from 100 m depth to the surface during the day. This depth range was chosen to sample through the depths known for paralarval *Ommastrephes bartramii*. Species with paralarval distributions deeper than this depth range will thus be under-represented. In total, 404 tows were conducted at 59 stations. All HM surveys used a 4 m² ring net equipped with 0.505 mm mesh, a General Oceanics flowmeter, and a Benthos time-depth recorder (TDR). The TC-91 survey also used a 4 m² ring net. The TC-92 survey used a 2 m² ring net for 94 tows and a 4 m² ring net for 4 tows. TC nets carried a Wildlife Computers TDR. Bower (1996) described the sampling procedure in detail.

Net collections were fixed at sea in 4 to 6% buffered formalin for 3 to 6 h, then preserved in 50% isopropyl alcohol. During the HM-91 survey, paralarvae were sorted from the plankton at sea with dissecting microscopes. Samples from subsequent TC and HM surveys were sorted on shore. Paralarvae were identified to genus and species level with the aid of published figures and descriptions (Harman & Young 1985, Young & Harman 1985, 1987, 1989, Young et al. 1985, Young & Hirota 1990, Young 1991). In some genera, more than 1 species was present, but identification was not possible either because a growth series connecting paralarvae to known adults was not available or the adults were known but undescribed. In such cases, species were given letter designations. Abundances of each species were standardized (individuals per 50 m² of sea surface sampled) for each tow (Smith & Richardson 1977). Station abundances were calculated as the average of all station tows.

Assemblage groups. Assemblage groups were determined with the 2-way indicator

species analysis program TWINSpan (Hill 1979, Gauch & Whittaker 1981) using PC-ORD software (McCune & Mefford 1997). Analyses are based on abundance data from all surveys for species with total catches greater than 15. Abundances at each station were used for comparisons of onshore-offshore distribution to determine which species preferentially spawn near the islands (Spearman rank test, $p < 0.05$).

RESULTS

Family and species composition

The 404 tows at 59 stations collected 10 375 paralarvae from 21 families and 58 species (Table 1). The most numerous families were the Ommastrephidae (23% of total catch), Pyroteuthidae (17%), Enoploteuthidae (16%), Onychoteuthidae (14%), and Ctenopterygidae (8%). The most numerous species were *Ommastrephes bartramii* (18%), *Pterygioteuthis microlampas* (15%), *Ctenopteryx sicula* (8%), and *Onychoteuthis compacta* (6%). Interannual catch rates for most species varied widely. Catches of the most abundant species showed a general decline over the 3 survey years. Results for the more frequently taken taxa are presented below.

Species distribution: squids

A total of 8848 squids from 16 families and 44 species were collected.




Fig. 1. Distribution and abundance of *Ommastrephes bartramii* paralarvae collected near the Hawaiian Islands during 5 cruises in 1991 to 1993. Circles represent mean station abundance (ind./50 m²). White circles: 1991 catches; gray circles: 1992 catches; black circles: 1993 catches. Circle area represents catch abundance. X: no catch

Family Ommastrephidae. *Ommastrephes bartramii* was the most numerous cephalopod species collected, composing 18% of the cephalopod catch and 80% of the ommastrephid catch; catches occurred at 41 stations (Fig. 1), and 40% were collected more than 200 km offshore. *Hyaloteuthis pelagica* was also widely distributed, occurring at 37 stations; 35% were collected more than 200 km offshore. Catches of the other 2 ommastrephids represented in the collections occurred farther inshore. *Sthenoteuthis oualaniensis* was collected at 26 stations (Fig. 2); 40% occurred less than 25 km offshore. *Nototodarus hawaiiensis* was collected at 15 stations; 54% occurred less than 25 km offshore.

Family Pyroteuthidae. All 3 pyroteuthid species were widely distributed. *Pterygioteuthis microlampas* composed 89% of the pyroteuthid catch and was the most commonly collected cephalopod species, occurring in 68% of the tows and at 56 stations (Fig. 3); 41% occurred more than 200 km offshore. *Pterygioteuthis giardi* was collected at 26 stations; 54% occurred more than 200 km offshore. *Pyroteuthis addolux* was collected at 17 stations; 81% occurred more than 200 km offshore.

Family Enoploteuthidae. *Abraliopsis* sp. A (Young & Harman 1985) was collected at 46 stations (Fig. 4) and composed 30% of the enoploteuthid catch; 66% occurred more than 200 km offshore. *Enoploteuthis*

Table 1. Cephalopod paralarvae collected near the Hawaiian Islands during 5 surveys in 1991 to 1993. Numbers for higher taxa include specimens that could not be further identified. HM: 'Hokusei Maru'; TC: 'Townsend Cromwell'. 91: 1991; 92: 1992; 93: 1993. Mean abundance: ind./50 m². NE: distance (km) from archipelago to northeasternmost point of capture; SW: distance (km) from archipelago to southwesternmost point of capture. This point will lie windward (northeast) of the archipelago if no captures were made south of the archipelago. W: windward of the archipelago; L: leeward of the archipelago. na: not applicable

	HM-91	TC-91	HM-92	TC-92	HM-93	Total	Positive tows (%)	Positive-tow mean abundance (SD)	Offshore range NE SW	% collected < 50 km offshore
Order Octopoda	29	333	82	17	128	589				
Family Argonautidae	2	3	3	2	4	14				
<i>Argonauta argo</i>	2	3	3	2	4	14	3	0.8 (0.4)	486-W 14-W	21
Family Bolitaenidae	1	1	6	0	9	17				
<i>Eledonella pygmaea</i>	1	1	6	0	9	17	3	0.9 (0.3)	736-W 204-W	0
Family Octopodidae	26	325	70	12	111	544				
<i>Octopus cyanea</i>	14	250	38	3	83	388	18	4.3 (7.9)	264-W 256-L	82
<i>Octopus ornatus</i>	0	2	3	0	0	5	1	1.0 (0.3)	26-W 9-W	100
<i>Octopus</i> Type A ^a	1	3	0	0	2	6	1	1.0 (0.6)	26-W 113-L	83
<i>Octopus</i> Type B ^a	0	13	2	4	7	26	3	2.0 (1.1)	143-W 113-L	73
<i>Octopus</i> Type C ^a	1	11	1	1	5	19	4	1.0 (0.5)	264-W 34-L	84
<i>Octopus</i> Type D ^a	1	1	1	1	0	4	1	1.2 (0.4)	59-W 126-L	50
<i>Octopus</i> Type E ^a	4	18	1	2	0	25	3	2.2 (2.4)	59-W 11-L	92
<i>Octopus</i> Type H ^a	2	14	8	1	13	38	7	1.0 (0.6)	351-W 113-L	66
<i>Octopus</i> Type I ^a	3	13	9	0	1	26	5	1.1 (0.6)	238-W 34-L	73
Family Tremoctopodidae	0	4	3	3	4	14				
<i>Tremoctopus violaceus</i>	0	4	3	3	4	14	3	1.1 (0.5)	762-W 215-L	14
Order Sepioidea	1	1	0	0	0	2				
Family Sepiolidae	1	1	0	0	0	2				
<i>Euprymna scolopes</i>	1	1	0	0	0	2	1	1.0 (0.1)	26-W 34-L	100
Order Teuthida	1790	1874	2263	1293	1628	8848				
Family Ancistrocheiridae	3	9	4	7	2	25				
<i>Ancistrocheirus lesueurii</i>	3	9	4	7	2	25	5	1.1 (0.8)	721-W 212-L	36
Family Brachioteuthidae	101	73	117	45	56	392				
<i>Brachioteuthis</i> sp.	101	73	117	45	56	392	42	2.1 (2.2)	1169-W 256-L	43
Family Chiroteuthidae	31	20	48	18	12	129				
<i>Chiroteuthis picteti</i>	3	10	5	0	3	21	4	1.3 (0.6)	183-W 113-L	57
<i>Chiroteuthis</i> sp. nov. ^b	27	7	43	16	8	101	17	1.2 (0.9)	762-W 126-L	28
<i>Grimalditeuthis bonplandi</i>	0	3	0	2	1	6	1	1.4 (0.6)	1169-W 113-L	50
<i>Plancoteuthis danae</i>	1	0	0	0	0	1	0.2	1.6 (na)	106-W 106-W	0
Family Ctenopterygidae	192	103	272	65	141	773				
<i>Ctenopteryx sicula</i>	192	103	272	65	141	773	59	2.7 (2.9)	1169-W 256-L	26

(Table continued overleaf)

Table 1 (continued)

	HM-91	TC-91	HC-92	TC-92	HM-93	Total	Positive tows (%)	Positive-tow mean abund- ance (SD)	Offshore range NE SW	% collected < 50 km offshore
Family Cranchiidae	96	214	209	32	188	739				
<i>Cranchia scabra</i>	29	28	95	7	69	228	34	1.2 (0.8)	736-W 256-L	29
<i>Galiteuthis pacifica</i>	1	0	5	1	5	12	3	0.7 (0.3)	1169-W 9-W	25
<i>Helicocranchia</i> sp. B ^c	40	59	15	4	52	170	14	3.0 (4.9)	979-W 207-L	43
<i>Leachia pacifica</i>	16	25	25	12	16	94	18	1.1 (0.7)	721-W 256-L	49
<i>Liocranchia reinhardtii</i>	6	67	50	4	25	152	22	1.3 (1.3)	471-W 256-L	70
<i>Liocranchia valdiviae</i>	4	33	18	4	19	78	14	1.2 (1.0)	721-W 207-L	62
<i>Megalocranchia fisheri</i>	0	2	1	0	2	5	1	0.8 (0.2)	230-W 30-L	60
Family Cycloteuthididae	0	1	0	0	0	1				
<i>Cycloteuthis sirventi</i>	0	1	0	0	0	1	0.2	0.8 (na)	14-W 14-W	100
Family Enoploteuthididae	357	351	416	121	270	1515				
<i>Abralia astrosticta</i>	0	6	0	0	1	7	1	1.7 (1.1)	71-W 14-W	86
<i>Abralia trigonura</i>	108	185	34	17	27	371	20	4.4 (5.1)	264-W 113-L	88
<i>Abraliopsis pacificus</i>	39	39	82	39	80	279	38	1.5 (1.2)	1169-W 256-L	50
<i>Abraliopsis</i> sp. A ^d	101	38	180	48	86	453	45	2.2 (2.3)	1169-W 256-L	24
<i>Enoploteuthis higginsii</i>	78	67	78	13	52	288	37	1.7 (1.6)	721-W 256-L	39
<i>Enoploteuthis jonesi</i>	1	6	8	0	6	21	4	1.0 (0.6)	264-W 207-L	81
<i>Enoploteuthis reticulata</i>	30	10	34	4	15	93	16	1.3 (1.4)	736-W 256-L	27
Family Histioteuthididae	8	26	7	5	8	54				
<i>Histioteuthis hoylei</i>	0	14	6	0	4	24	3	1.6 (1.2)	143-W 207-L	79
<i>Histioteuthis</i> sp. B ^c	0	3	0	0	1	4	1	1.8 (1.6)	230-W 204-W	0
<i>Histioteuthis</i> sp. C ^c	7	9	1	1	3	21	4	1.2 (1.0)	581-W 113-L	33
<i>Histioteuthis</i> sp. D ^c	1	0	0	4	0	5	1	1.3 (0.3)	979-W 351-W	0
Family Lepidoteuthididae	2	0	0	1	0	3				
<i>Lepidoteuthis grimaldii</i>	2	0	0	1	0	3	1	1.2 (0.3)	955-W 16-W	33
Family Mastigoteuthididae	0	5	1	0	0	6				
<i>Mastigoteuthis famelica</i>	0	5	1	0	0	6	1	0.9 (0.2)	183-W 14-W	83
Family Octopoteuthididae	10	9	10	9	12	50				
<i>Octopoteuthis nielsenii</i>	10	9	10	9	8	46	9	1.2 (0.7)	1169-W 207-L	30
<i>Taningia danae</i>	0	0	0	0	4	4	1	1.0 (0.2)	30-L 207-L	50
Family Ommastrephidae	403	210	378	634	519	2144				
<i>Hyaloteuthis pelagica</i>	22	65	90	34	40	251	27	2.1 (4.0)	736-W 256-L	49
<i>Nototodarus hawaiiensis</i>	6	11	11	1	6	35	6	1.0 (0.4)	264-W 113-L	80
<i>Ommastrephes bartramii</i>	368	100	242	556	453	1719	47	8.1 (11.9)	721-W 256-L	34
<i>Sthenoteuthis oualaniensis</i>	7	34	35	43	20	139	19	1.9 (2.2)	581-W 256-L	63
Family Onychoteuthididae	156	444	343	103	251	1297				
<i>Onychoteuthis compacta</i>	79	226	114	26	93	538	46	2.4 (3.3)	1169-W 256-L	33
<i>Onychoteuthis</i> sp. B ^c	0	3	4	0	7	14	3	0.8 (0.3)	264-W 256-L	64
<i>Onychoteuthis</i> sp. C ^c	34	166	117	13	87	417	36	2.3 (2.3)	721-W 256-L	73
<i>Onykia</i> sp. A ^c (<i>O. carriboea</i> ?)	38	47	103	61	58	307	43	1.5 (1.1)	721-W 256-L	36
<i>Onykia</i> sp. B ^c (<i>O. rancureli</i> ?)	5	1	5	3	0	14	3	1.5 (0.7)	979-W 16-W	29
Family Pholidoteuthididae	10	11	0	1	2	24				
<i>Pholidoteuthis</i> sp.	10	11	0	1	2	24	3	1.7 (1.0)	721-W 22-W	8
Family Pyroteuthididae	411	377	419	234	147	1588				
<i>Pterygoteuthis giardi</i>	16	40	35	1	15	107	13	1.8 (1.2)	955-W 215-L	45
<i>Pterygoteuthis microlampas</i>	373	331	360	227	125	1416	68	5.0 (6.2)	955-W 256-L	40
<i>Pyroteuthis addolux</i>	22	6	21	6	7	62	8	2.2 (3.0)	979-W 215-L	8
Family Thysanoteuthididae	10	21	39	18	20	108				
<i>Thysanoteuthis rhombus</i>	10	21	39	18	20	108	18	1.2 (0.7)	721-W 256-L	42
Unidentified	185	180	371	133	67	936				
Total cephalopods	2005	2388	2716	1443	1823	10375				
Total species	45	52	46	41	47	57				

^aYoung & Harman (1989); ^bYoung (1991); ^cidentified from growth series; ^dYoung & Harman (1985); ^eYoung & Harman (1987)

higginsii, *Abraliopsis pacificus* and *E. reticulata* also had wide latitudinal distributions. *Abralia trigonura* (Fig. 5) and *E. jonesi* were collected much closer to shore; 40% and 67% of their respective catches occurred less than 25 km offshore.

Family Onychoteuthidae. *Onychoteuthis compacta* was collected at 50 stations (Fig. 6) and composed 43% of the onychoteuthid catch; 59% occurred more than 200 km offshore. *Onykia* sp. A was also widely distributed at 50 stations, but less abundant than *O. compacta*; 40% occurred more than 200 km offshore. *Onychoteuthis* sp. C (Young & Harman 1987) was widely distributed at 41 stations (Fig. 7), yet 53% occurred less than 25 km offshore. *Onychoteuthis* sp. B (Young & Harman 1987) was collected at 9 stations; 50% occurred less than 25 km offshore.

Family Ctenopterygidae. *Ctenopteryx sicula* was the most widely distributed cephalopod collected, occurring at 97% of the stations (Fig. 8); 57% were collected more than 200 km offshore.

Family Cranchiidae. *Cranchia scabra* was collected at 42 stations (Fig. 9) and composed 31% of the cranchiid catch; 58% occurred more than 200 km offshore. *Helicocranchia* sp. B was collected at 22 stations and composed 23% of the cranchiid catch; 50% occurred more than 200 km offshore. *Leachia pacifica* was also widely distributed; catches occurred at 38 stations, and 37% were collected more than 200 km offshore. *Liocranchia reinhardtii* and *Liocranchia valdiviae* occurred closer to shore; 46 and 41% of their respective catches occurred less than 25 km offshore.

Species distribution: octopods

A total of 589 octopods from 4 families and 12 species were collected. The 9 octopodid species composed 92% of the octopodid numeric catch. All octopodids were collected less than 352 km from the islands; 59% occurred less than 25 km offshore. *Octopus cyanea* was collected at 23 stations (Fig. 10) and composed 72% of the octopodid numeric catch. *Octopus* Type H (Young & Harman 1989) had the furthest offshore distribution of the octopodids; 29% were collected more than 200 km off-




Fig. 2. Distribution and abundance of *Sthenoteuthis oualaniensis*. Symbols described in Fig. 1




Fig. 3. Distribution and abundance of *Pterygoteuthis microlampus*. Symbols described in Fig. 1




Fig. 4. Distribution and abundance of *Abraliopsis* sp. A. Symbols described in Fig. 1




Fig. 5. Distribution and abundance of *Abralia trigonura*. Symbols described in Fig. 1




Fig. 6. Distribution and abundance of *Onychoteuthis compacta*. Symbols described in Fig. 1




Fig. 7. Distribution and abundance of *Onychoteuthis* sp. C. Symbols described in Fig. 1

shore. *Eledonella pygmaea* was the only cephalopod collected exclusively offshore; catches occurred at 8 windward stations north of 25°02' N greater than 200 km offshore. *Tremoctopus violaceus* was collected at 11 stations and had the widest latitudinal range of all octopods (21°20' to 29°14' N); 79% occurred more than 200 km offshore.

Assemblages

Assemblage analysis using the 2-way indicator species analysis program defined 2 distributional assemblages: 'island associated' and 'oceanic' (Table 2). The 15 'island-associated' species include 6 species with benthic adults (*Octopus* spp.), 6 species with mesopelagic adults (*Abralia trigonura*, *Liocranchia reinhardtii*, *L. valdiviae*, *Histioteuthis hoylei*, *Chroteuthis picteti*, and *Enoploteuthis jonesi*) and 3 species with epipelagic adults (*Onychoteuthis* sp. C, *Sthenoteuthis oualaniensis* and *Nototodarus hawaiiensis*). Analysis of onshore-offshore abundance patterns showed that all 'island-associated' species decreased significantly in abundance with increasing distance offshore and had more than 50% of their paralarvae collected within 50 km of shore.

DISCUSSION

Paralarval distribution patterns provide further evidence that *Abralia trigonura*, *Liocranchia reinhardtii* and *Chroteuthis picteti* are members of the Hawaiian Mesopelagic Boundary Community (MBC). However, the present data lend no support to Young's (1995) suggestion that *Pterygioteuthis giardi* is a probable facultative boundary species.

Paralarval catch data indicate that *Liocranchia valdiviae*, *Histioteuthis hoylei*, and *Enoploteuthis jonesi* are also probable MBC members. *L. valdiviae* adults were rarely caught near the Hawaiian Islands during MBC studies; however, the subadults of this deep-living species, unlike those of their congener, do not vertically migrate (Young 1978) and were thus mostly out of the sampling range of those studies. *H. hoylei* is widely distributed in

the Pacific (Voss et al. 1998), but appears to spawn preferentially near land masses. Mature *H. hoylei* females collected off Japan occur in aggregations near an isolated oceanic rise, suggesting this species might spawn near the seafloor, as reported for *H. miranda* and *H. celetaria pacifica* (Nesis 1993a). *H. reversa* paralarvae also commonly occur near land masses (Clarke 1966). *E. jonesi* occurs in Hawaiian and equatorial waters (Burgess 1982) and has not been considered a member of the Hawaiian MBC (Young 1995); however, the present study indicates that this species is a nearshore spawner. Some other *Enoplo-teuthis* species have been reported to spawn only over the slopes or in nearshore oceanic regions (Nesis 1993a, 1996).

The paralarvae of *Sthenoteuthis oualaniensis*, an abundant member of the oceanic nekton throughout the tropical and subtropical Indo-Pacific region (Roper et al. 1984, Young & Hirota 1998), were also 'island associated'. Increased catches of *S. oualaniensis* paralarvae in nearshore waters also have been reported from the Saya de Malha Bank, Seychelles, and its slopes (Nesis 1993b) and the Arabian Sea (Piatkowski et al. 1993). Most mature *S. oualaniensis* females collected in the present study area occur along the windward slope of the islands (Suzuki et al. 1986, Young & Hirota 1998), further suggesting that nearshore waters are favorable for spawning.

Some epipelagic squids near the Hawaiian Islands, including *Sthenoteuthis oualaniensis* and possibly *Onychoteuthis* sp. C, may migrate inshore before spawning. Nesis (1993a) has suggested that other oceanic squids (*Todarodes sagittatus*, *Ornithoteuthis* spp. and *Pholidoteuthis* spp.) similarly migrate to seamounts to spawn (Nesis 1993a). Inshore-offshore spawning migrations are common in the Loliginidae (Mangold 1987); these squid must find a solid substrate where they can attach their egg capsules.

Possibly good feeding conditions near the Hawaiian Islands could explain why some oceanic cephalopods appear to spawn preferentially nearshore. Young pelagic squids can consume 80 to 100% of their body weight per day (LaRoe 1971, Hurley 1976). Meeting such a high food requirement may be difficult, particularly in oligotrophic




Fig. 8. Distribution and abundance of *Ctenopteryx sicula*. Symbols described in Fig. 1




Fig. 9. Distribution and abundance of *Cranchia scabra*. Symbols described in Fig. 1




Fig. 10. Distribution and abundance of *Octopus cyanea*. Symbols described in Fig. 1

Table 2. Paralarval assemblages identified from 5 plankton surveys near the Hawaiian Islands. Species groups within assemblages were determined with a 2-way synthesis table technique (TWINSPAN). Only species with ≥ 15 paralarvae collected were considered in the analyses. Literature references for species with letter designations are listed in Table 1

Island-associated assemblage	Oceanic assemblage
<i>Abralia trigonura</i>	<i>Abraliopsis pacificus</i>
<i>Chiroteuthis picteti</i>	<i>Abraliopsis</i> sp. A
<i>Enoploteuthis jonesi</i>	<i>Ancistrocheirus lesueurii</i>
<i>Histioteuthis hoylei</i>	<i>Brachioteuthis</i> sp.
<i>Liocranchia reinhardti</i>	<i>Chiroteuthis</i> sp. nov.
<i>Liocranchia valdiviae</i>	<i>Cranchia scabra</i>
<i>Nototodarus hawaiiensis</i>	<i>Ctenopteryx sicula</i>
<i>Octopus cyanea</i>	<i>Eledonella pygmaea</i>
<i>Octopus</i> Type B	<i>Enoploteuthis higginsii</i>
<i>Octopus</i> Type C	<i>Enoploteuthis reticulata</i>
<i>Octopus</i> Type E	<i>Helicocranchia</i> sp. B
<i>Octopus</i> Type H	<i>Histioteuthis</i> sp. C
<i>Octopus</i> Type I	<i>Hyaloteuthis pelagica</i>
<i>Onychoteuthis</i> sp. C	<i>Leachia pacifica</i>
<i>Sthenoteuthis oualaniensis</i>	<i>Octopoteuthis nielsenii</i>
	<i>Ommastrephes bartramii</i>
	<i>Onychoteuthis compacta</i>
	<i>Onykia</i> sp. A
	<i>Pholidoteuthis</i> sp.
	<i>Pterygioteuthis giardi</i>
	<i>Pterygioteuthis microlampas</i>
	<i>Pyroteuthis addolux</i>
	<i>Thysanoteuthis rhombus</i>

oceanic environments. The islands and seamounts that form the crest of the Hawaiian Ridge result in a number of mesoscale features, including seamount-induced biological and nutrient concentration disturbances (Boehlert & Genin 1987), which could result in advantageous feeding conditions near the islands. Higher productivity and plankton biomass levels very near the Hawaiian Islands (Gilmartin & Revelante 1974) may enhance paralarval survival for some species (see Dandonneau & Charpy 1985). Unfortunately specific conditions that would provide a superior paralarval nursery ground are not known for these waters.

Waters near the Hawaiian Islands also may serve as favorable feeding grounds for spawning adults. Boehlert & Mundy (1994) have suggested that this may be the case for spawning tuna. Pelagic species such as tuna and squid typically aggregate in waters near islands and banks (Uda & Ishino 1958) and above seamounts (Inoue 1983, Yasui 1986) to feed. Increasing evidence indicates that some cephalopods, including *Sthenoteuthis oualaniensis* (Harman et al. 1989, Nigmatullin & Laptikhovskiy 1994), are multiple (i.e. iteroparous *sensu lato*) rather than single/terminal (i.e. semelparous *sensu lato*) spawners. Mature females spawning multiple batches of eggs on spawning grounds presumably must feed to survive and reproduce again.

In conclusion, the paralarval fauna near the Hawaiian Islands is the most diverse yet reported (Table 3). Species that showed evidence of increased spawning near the islands include not only those with bottom or nearshore spawning, such as *Abralia trigonura*, *Nototodarus hawaiiensis*, and *Octopus* spp., but also oceanic species not connected with the bottom of nearshore areas, such as *Sthenoteuthis oualaniensis*. Improved feeding conditions in waters near the archipelago might be one reason for the apparent increase in nearshore spawning by some oceanic species.

Acknowledgements. This study was supported by grants from the East-West Center (Honolulu) and the Japan Society for the Promotion of Science (Tokyo) awarded to J.R.B., from Hawaii Sea Grant (grant # NA89AA-D-SG063) awarded to R.E.Y., and from the Joint Institute for Marine and Atmospheric Research (JIMAR) awarded to P.F. We appreciate reviews of the early manuscript by George Boehlert, Chung-Cheng Lu, Kir Nesis, Michael Vecchione and 2 anonymous reviewers. We also acknowledge the help of June Firing, Ilgiz Irnazarow, Takashi Okutani, Uwe Piatkowski, Stewart B. Reid and the officers and crews of the 'Hokusei Maru' and 'Townsend Cromwell'. We dedicate this paper to the memory of Kiyohiko Ishii.

LITERATURE CITED

- Arkhipkin AI, Zheronkin YN, Loktionov YA, Shchetinnikov AS (1988) Fauna and distribution of pelagic cephalopods larvae in the Gulf of Guinea. Zool Zh 67(10):1459–1467 (in Russian with English abstract)
- Berry SS (1914) The Cephalopoda of the Hawaiian Islands. Bull Bureau Fish 32(789):255–361
- Boehlert GW, Genin A (1987) A review of the effects of seamounts on biological processes. In: Keating BH, Fryer P, Batiza R, Boehlert GW (eds) Seamounts, islands and atolls. Geophys Monogr Am Geophys Union 43:319–334
- Boehlert GW, Mundy BC (1994) Vertical and onshore-off-shore distributional patterns of tuna larvae in relation to physical habitat features. Mar Ecol Prog Ser 107:1–13
- Bower JR (1994) Distribution of paralarvae of the squid *Ommastrephes bartramii* near the Hawaiian Archipelago. MSc thesis, Dept of Oceanography, University of Hawaii, Honolulu
- Bower JR (1996) Estimated paralarval drift and inferred hatching sites for *Ommastrephes bartramii* (Cephalopoda: Ommastrephidae) near the Hawaiian Archipelago. Fish Bull 94(3):398–411
- Boyle PR (1990) Cephalopod biology in the fisheries context. Fish Res 8:303–321
- Burgess LA (1982) Four new species of squid (Oegopsida: *Enoploteuthis*) from the Central Pacific and a description of adult *Enoploteuthis reticulata*. Fish Bull 80(4):703–734
- Clarke MR (1966) A review of the systematics and ecology of oceanic squids. Adv Mar Biol 4:91–300
- Dandonneau Y, Charpy L (1985) An empirical approach to the island mass effect in the south tropical Pacific based on sea surface chlorophyll concentrations. Deep-Sea Res 32: 707–721
- Dawe FG, Stephen SJ (1988) The cephalopod assemblage of the Gulf Stream System east of 60°W. Malacologia 29(1): 235–245

Table 3. Numbers of families and species of cephalopod paralarvae collected from different regions. *Octopods not reported

	Hawaiian Islands ^a	Pacific coast of Japan ^{b,c,d,e,f}	Gulf Stream east of 60° W ^g	Gulf of Guinea ^h	Northwest Pacific ^{i,j,k,l,m}	Arabian Sea ^{n,o,p}	California Current ^q
Order Octopoda							
Alloposidae	0		1	0	0	0	
Argonautidae	1		1	1	0	0	
Bolitaenidae	1		0	0	1	1	
Octopodidae	9		2	3	1	1	
Ocythoidae	0		0	1	0	0	
Tremoctopodidae	1		0	0	0	0	
Order Sepioidea							
Idiosepiidae	0	1	0	0	0	0	0
Sepiolidae	1	3	0	0	0	0	0
Order Teuthida							
Ancistrocheiridae	1	1	0	1	0	0	0
Bathyteuthididae	0	0	0	0	0	1	0
Brachyteuthididae	1	1	1	1	1	0	0
Chiroteuthididae	4	1	1	0	0	1	1
Chtenopterygidae	1	1	1	1	0	1	1
Cranchiidae	7	7	3	3	1	2	4
Cycloteuthididae	1	0	0	1	0	0	0
Enoploteuthididae	7	4	3	2	2	3	1
Gonatidae	0	2	2	0	9	0	1
Histioteuthididae	4	0	1	1	1	0	1
Joubiniteuthididae	0	0	1	0	0	1	0
Lepidoteuthididae	1	0	0	0	0	0	0
Loliginidae	0	2	0	0	0	0	1
Lycoteuthididae	0	0	1	0	0	0	0
Mastigoteuthididae	1	0	1	1	0	0	0
Neoteuthididae	0	0	0	0	0	0	0
Octopoteuthididae	2	1	1	0	1	1	1
Ommastrephidae	4	6	1	2	1	2	1
Onychoteuthididae	5	3	2	3	1	1	1
Pholidoteuthididae	1	0	0	0	0	0	0
Psychroteuthididae	0	0	0	0	0	0	0
Pyroteuthididae	3	1	4	2	0	0	1
Thysanoteuthididae	1	1	0	1	1	0	0
No. of families	21	15*	17	15	11	11	11*
No. of species	57	35*	27	24	20	15	14*
References: ^a Present study; ^b Okutani (1968); ^c Okutani (1969); ^d Sato & Sawada (1974); ^e Yamamoto & Okutani (1975); ^f Saito & Kubodera (1993); ^g Dawe & Stephen (1988); ^h Arkhipkin et al. (1988); ⁱ Okutani (1966); ^j Kubodera & Okutani (1981); ^k Kubodera & Jefferts (1984a); ^l Kubodera & Jefferts (1984b); ^m Kubodera (1991); ⁿ Nesis (1974); ^o Piatkowski & Welsch (1991); ^p Piatkowski et al. (1993); ^q Okutani & McGowan (1969)							

Gauch HG Jr, Whittaker RH (1981) Hierarchical classification of community data. *J Ecol* 69(2):537–557

Gilmartin M, Revelante N (1974) The 'island mass effect' on the phytoplankton and primary production of the Hawaiian Islands. *J Exp Mar Biol Ecol* 16:181–204

Harman RF, Young RE (1985) The larvae of ommastrephid squids (Cephalopoda, Teuthoidea) from Hawaiian waters. *Vie Milieu* 35(3/4):211–222

Harman RF, Young RE, Reid SB, Mangold KM, Suzuki T, Hixon RF (1989) Evidence for multiple spawning in the tropical oceanic squid *Stenoteuthis oualaniensis* (Teuthoidea: Ommastrephidae). *Mar Biol* 101:513–519

Hill MO (1979) TWINSPLAN—a FORTRAN program for arranging multivariate data in an ordered two-way table

by classification of the individuals and attributes. *Ecology and Systematics*, Cornell University, Ithaca, NY

Hurley AC (1976) Feeding behavior, food consumption, growth, and respiration of the squid *Loligo opalescens* raised in the laboratory. *Fish Bull* 74(1):176–182

Inoue M (1983) Exploitation of fishing grounds of southward migrants of fall skipjacks in the northwestern Pacific Ocean. *J Fac Mar Sci Technol Tokai Univ* 17:121–130 (in Japanese with English abstract)

Kubodera T (1991) Distribution and abundance of the early life stages of octopus, *Octopus dofleini* Walker 1910 in the North Pacific. *Bull Mar Sci* 49(1–2):235–243

Kubodera T, Jefferts K (1984a) Distribution and abundance of the early life stages of squid, primarily Gonatidae (Ceph-

- alopoda, Oegopsida), in the northern North Pacific (Part 1). Bull Natl Sci Mus Ser A (Zool) 10(3):91–106
- Kubodera T, Jefferts K (1984b) Distribution and abundance of the early life stages of squid, primarily Gonatidae (Cephalopoda, Oegopsida), in the northern North Pacific (Part 2). Bull Natl Sci Mus Ser A (Zool) 10(4):165–193
- Kubodera T, Okutani T (1981) The systematics and identification of larval cephalopods from the northern North Pacific. Res Inst N Pac Fish Hokkaido Univ Spec Vol, p 131–159
- LaRoe ET (1971) The culture and maintenance of the loliginid squids *Sepioteuthis sepioidea* and *Doryteuthis plei*. Mar Biol 9(1):9–25
- Mangold K (1987) Reproduction. In: Boyle PR (ed) Cephalopod life cycles, Vol 2. Academic Press Inc, London, p 157–200
- McCune B, Mefford MJ (1997) PC-ORD. Multivariate analysis of ecological data, Version 3.0. MjM Software Design, Gleneden Beach, OR
- Nesis KN (1974) Cephalopod larvae in the western Arabian Sea. Oceanology 14(3):441–445
- Nesis K (1993a) Cephalopods of seamounts and submarine ridges. In: Okutani T, O'Dor RK, Kubodera T (eds) Recent advances in cephalopod fisheries biology. Tokai University Press, Tokyo, p 365–373
- Nesis K (1993b) Population structure of oceanic ommastrephids, with particular reference to *Sthenoteuthis oualaniensis*: a review. In: Okutani T, O'Dor RK, Kubodera T (eds) Recent advances in cephalopod fisheries biology. Tokai University Press, Tokyo, p 375–383
- Nesis KN (1996) Mating, spawning, and death in oceanic cephalopods: a review. Ruthenica 6(1):23–64
- Nigmatullin CM, Laptikhovskiy VV (1994) Reproductive strategies in the squids of the family Ommastrephidae (preliminary report). Ruthenica 4(1):79–82
- Okutani T (1966) Studies on early life history of decapodan Mollusca—II. Planktonic larvae of decapodan cephalopods from the northern North Pacific in summer seasons during 1952–1959. Bull Tokai Reg Fish Res Lab 45:61–79
- Okutani T (1968) Studies on early life history of decapodan Mollusca—III. Systematics and distribution of larvae of decapod cephalopods collected from the sea surface on the Pacific coast of Japan 1960–1965. Bull Tokai Reg Fish Res Lab 55:9–57
- Okutani T (1969) Studies on early life history of decapodan Mollusca—IV. Bull Tokai Reg Fish Res Lab 58:83–96
- Okutani T, McGowan JA (1969) Systematics, distribution, and abundance of the euplanktonic squid (Cephalopoda, Decapoda) larvae of the California Current April 1954 – March 1957. Bull Scripps Inst Oceanogr Univ Calif 14:1–90
- Piatkowski U, Welsch W (1991) On the distribution of pelagic cephalopods in the Arabian Sea. Bull Mar Sci 49(1-2): 186–198
- Piatkowski U, Welsch W, Ropke A (1993) Distribution patterns of the early life stages of pelagic cephalopods in three geographically different regions of the Arabian Sea. In: Okutani T, O'Dor RK, Kubodera T (eds) Recent advances in cephalopod fisheries biology. Tokai University Press, Tokyo, p 417–431
- Reid SB, Hirota J, Young RE, Hallacher LE (1991) Mesopelagic-boundary community in Hawaii: micronekton at the interface between neritic and oceanic systems. Mar Biol 109:427–440
- Roper CFE, Sweeney MJ, Nauen CE (1984) FAO species catalogue. Vol 3. An annotated and illustrated catalogue of species of interest to fisheries. Cephalopods of the world. FAO Fish Synop 125, FAO, Rome
- Saito H, Kubodera T (1993) Distribution of ommastrephid rhynchoteuthion paralarvae (Mollusca, Cephalopoda) in the Kuroshio region. In: Okutani T, O'Dor RK, Kubodera T (eds) Recent advances in cephalopod fisheries biology. Tokai University Press, Tokyo, p 457–466
- Sato K, Sawada T (1974) Studies on larvae of decapod cephalopods in the waters of the Izu Peninsula—I. Identification of larvae. Bull Shizuoka Prefect Fish Exp Stn 8: 11–21 (in Japanese)
- Smith PE, Richardson SL (1977) Standard techniques for pelagic fish egg and larva surveys. FAO Fish Tech Pap 175, FAO, Rome
- Suzuki T, Yamamoto S, Ishii K, Matsumoto WM (1986) On the flying squid *Stenoteuthis* [sic] *oualaniensis* (Lesson) in Hawaiian waters. Bull Fac Fish Hokkaido Univ 37(2):111–123
- Uda M, Ishino M (1958) Enrichment pattern resulting from eddy systems in relation to fishing grounds. J Tokyo Univ Fish 44:105–129
- Vecchione M (1987) Juvenile ecology. In: Boyle PR (ed) Cephalopod life cycles, Vol 2. Academic Press Inc, London, p 61–84
- Voss NA, Nesis KN, Rodhouse PG (1998) The cephalopod family Histioteuthidae (Oegopsida): systematics, biology, and biogeography. Smithsonian Contrib Zool 586:293–372
- Yamamoto K, Okutani T (1975) Studies on early life history of decapodan Mollusca—V. Systematics and distribution of epipelagic larvae of decapod cephalopods in the south-western waters of Japan during the summer in 1970. Bull Tokai Reg Fish Res Lab 83:45–96
- Yasui M (1986) Albacore, *Thunnus alalunga*, pole-and-line fishery around the Emperor [sic] Seamounts. In: Uchida RN, Hayasi S, Boehlert GW (eds) Environment and resources of seamounts in the North Pacific. NOAA Tech Rep NMFS 43:37–40
- Young RE (1978) Vertical distribution and photosensitive vesicles of pelagic cephalopods from Hawaiian waters. Fish Bull 76(3):583–615
- Young RE (1991) Chiroteuthid and related paralarvae from Hawaiian waters. Bull Mar Sci 49(1-2):162–185
- Young RE (1995) Aspects of the natural history of pelagic cephalopods of the Hawaiian mesopelagic-boundary region. Pac Sci 49(2):143–155
- Young RE, Harman RF (1985) Early life history stages of enoploteuthin squids (Cephalopoda: Teuthoidea: Enoploteuthidae) from Hawaiian waters. Vie Milieu 35(3/4):181–201
- Young RE, Harman RF (1987) Descriptions of the larvae of three species of the *Onychoteuthis banksii* complex from Hawaiian waters. Veliger 29(3):313–321
- Young RE, Harman RF (1989) Octopodid paralarvae from Hawaiian waters. Veliger 32(2):152–165
- Young RE, Hirota J (1990) Description of *Ommastrephes bartramii* (Cephalopoda: Ommastrephidae) paralarvae with evidence of spawning in Hawaiian waters. Pac Sci 44(1): 71–80
- Young RE, Hirota J (1998) Review of the ecology of *Sthenoteuthis oualaniensis* near the Hawaiian Archipelago. In: Okutani T (ed) Contributed papers to international symposium on large pelagic squids. Japan Marine Fishery Resources Research Center, Tokyo, p 131–143
- Young RE, Harman RF, Mangold KM (1985) The eggs and larvae of *Brachioteuthis* sp. (Cephalopoda: Teuthoidea) from Hawaiian waters. Vie Milieu 35(3/4):203–209